Epigenetic aging differ in breast cancer discordant twin pairs - Same life but different disease status.

Hannes Bode¹, Aino Hekkinen¹, Sara Lundgren¹, Arina Tagmazian¹, Miina Ollikainen¹, Jaakko Kaprio¹ ¹FIMM, HiLIFE, University of Helsinki; **Contact: <u>hannes.bode@helsinki.fi</u>** Visit:https://www.cancerprev.com/ or https://www2.helsinki.fi/en/hilife-fimm







MATERIAL AND METHODS

DNA methylation was quantified from 24 breast cancer discordant all female twin pairs, 11 monozygotic and 13 dizygotic, from the Finnish twin cohort [Kaprio, 2019]. These twin pairs were age 38 to 82 at the time point of sampling. A total of 10 pairs were sampled up to 15 years before the diagnosis, 14 pairs up to 30 years after the diagnosis. DNA methylation based epigenetic age and its deviation from chronological age, called epigenetic age acceleration (EAA) were calculated based on the Horvath [Horvath, 2013] and Phenoage [Levine, 2018] models.

Paired t-tests were applied to investigate differences in EAA between the healthy and cancer diagnosed twin in each pair. This was done separately for twin-pairs sampled before (Fig. 3 and Fig. 4, left), and after the diagnosis (Fig. 3 and Fig. 4, right).

Linear regression was used to assess any within-pair differences in EAA, and whether that relates to the time of diagnosis (Fig. 1 and Fig.2).

RESULTS



Fig.1 Linear regression of intra-pair differences in EAA (Horvath) by time to diagnosis, including the 95% confident band.



Fig.3 Plot of EAA (Horvath) of twin samples taken before and after the diagnosis The p-values and dEEA refer to the within-pair differences from a paired t-test.

Discussion

Breast cancer discordant twin pairs sampled before the diagnosis **do not show differences** in their epigenetic age acceleration. (Fig. 3 and Fig. 4, left)



<u>Fig.2</u> Linear regression of intra-pair differences in EAA (Phenoage) by time to diagnosis, including the 95% confident band.



<u>Fig.4</u> Plot of EAA (Phenoage) of twin samples taken before and after the diagnosis The p-values and dEEA refer to the within-pair differences from a paired t-test.

References:

Kaprio J, Bollepalli S, Buchwald J, Iso-Markku P,

Breast cancer discordant twin pairs sampled **after** the diagnosis **show differences** in their epigenetic age acceleration. (Fig. 3 and Fig. 4, right)

- The breast cancer diagnosed twin seem to undergo faster epigenetic aging.
- However, EEA is only significantly different after the time of diagnosis, and epigenetic alterations related to cancer treatment could explain the accelerated aging rate.

The intra-pair difference in EAA is larger after the diagnosis compared with before the diagnosis. (Fig. 1 and Fig. 2) However, longitudinal data is needed to confirm if the difference in EAA increases linearly over time.

• A possible underlying effect of breast cancer might be already present shortly before the diagnosis.

An additional number of samples of about 100 breast cancer discordant twin pairs are currently measured for their blood DNA methylation levels, to increase the cohort and enable further analysis.

Korhonen T, Kovanen V, Kujala U, Laakkonen EK, Latvala A, Leskinen T, Lindgren N, Ollikainen M, Piirtola M, Rantanen T, Rinne J, Rose RJ, Sillanpää E, Silventoinen K, Sipilä S, Viljanen A, Vuoksimaa E, Waller K. The Older Finnish Twin Cohort - 45 Years of Follow-up. Twin Res Hum Genet. 2019 Aug;22(4):240-254. doi: 10.1017/thg.2019.54. Epub 2019 Aug 29. PMID: 31462340.

Horvath S. DNA methylation age of human tissues and cell types [published correction appears in Genome Biol. 2015;16:96]. Genome Biol. **2013**;14(10):R115. doi:10.1186/gb-2013-14-10-r115

Levine ME, Lu AT, Quach A, et al. An epigenetic biomarker of aging for lifespan and healthspan. Aging (Albany NY). 2018;10(4):573-591. doi:10.18632/aging.101414

FIMM - Institute for Molecular Medicine Finland P.O. Box 20, FI-00014 University of Helsinki, Finland Biomedicum Helsinki 2U, Tukholmankatu 8, 00290 Helsinki, Finland

www.fimm.fi

In close partnership with:

